



INYO COUNTY PUBLIC HEALTH BRIEF

A Division of Health & Human Services
Richard O. Johnson, M.D., MPH
Public Health Officer, Inyo County
760-914-0496
drrickjohn@gmail.com



Public Health
Prevent. Promote. Protect.

2009 H1N1 Influenza Vaccine and Pregnant Women

September 22, 2009

Q: Why does CDC recommend that pregnant women receive the 2009 H1N1 influenza vaccine?

A. It is important for a pregnant woman to receive the 2009 H1N1 influenza vaccine as well as a seasonal influenza vaccine. A pregnant woman who gets any type of flu is at risk for serious complications and hospitalization. Pregnant women who are otherwise healthy have been severely impacted by the 2009 H1N1 influenza virus (formerly called “novel H1N1 flu” or “swine flu”). In comparison to the general population, a greater proportion of pregnant women infected with the 2009 H1N1 influenza virus have been hospitalized. In addition, severe illness and death has occurred in pregnant women. Six percent of confirmed fatal 2009 H1N1 flu cases thus far have been in pregnant women while only about 1% of the general population is pregnant. While hand washing, staying away from ill people, and other steps can help to protect pregnant women from influenza, vaccination is the single best way to protect against the flu.

Q: Is there a particular kind of flu vaccine that pregnant women should get? Are there flu vaccines that pregnant women should not get?

A. There are two type of flu vaccine. Pregnant women should get the “flu shot”— an inactivated vaccine (containing fragments of killed influenza virus) that is given with a needle, usually in the arm. The flu shot is approved for use in pregnant women.

The other type of flu vaccine — nasal-spray flu vaccine (sometimes called LAIV for “live attenuated influenza vaccine)—is not currently approved for use in pregnant women. This vaccine is made with live, weakened flu viruses that do not cause the flu). LAIV (FluMist®) is approved for use in healthy people 2-49 years of age who are not pregnant.

Q. Will the seasonal flu vaccine also protect against the 2009 H1N1 flu?

A. The seasonal flu vaccine is not expected to protect against the 2009 H1N1 flu. Similarly, the 2009 H1N1 influenza vaccine will not protect against seasonal influenza.

Q. Can the seasonal influenza vaccine and the 2009 H1N1 influenza vaccine be given at the same time?

A. Seasonal flu and 2009 H1N1 vaccines may be administered on the same day but given at different sites (e.g. one shot in the left arm and the other shot in the right arm). However, the seasonal vaccine is available now, but the 2009 H1N1 influenza vaccine will not be available until later in October. The usual seasonal influenza viruses are still expected to cause illness this fall and winter. Pregnant women and others at increased risk of complications of influenza are encouraged to get their seasonal flu vaccine NOW!

Q: Is the 2009 H1N1 influenza vaccine safe for pregnant women?

A: Influenza vaccines have not been shown to cause harm to a pregnant woman or her baby. The seasonal flu shot (injection) is proven as safe and already recommended for pregnant women in all trimesters. The 2009 H1N1 influenza vaccine is made using the same processes and facilities that are used to make seasonal influenza vaccines.

Q: What safety studies have been done on the 2009 H1N1 influenza vaccine and have any been done in pregnant women?

A: A number of clinical trials which test 2009 H1N1 influenza vaccine in healthy children and adults are underway. These studies are being conducted by the National Institutes of Allergies and Infectious Diseases (NIAID). Studies of 2009 H1N1 influenza vaccine in pregnant women are expected to be completed in September, and results shared in October.

Q: Does the 2009 H1N1 influenza vaccine have preservative in it?

A: In California, all vaccines given to pregnant women are thimerosal free. Manufacturers are producing preservative-free seasonal and 2009 H1N1 influenza vaccines in single dose syringes for pregnant women and small children.

Q. How many doses of the 2009 H1N1 flu vaccine will pregnant women need to get?

A. The U.S. Food and Drug Administration (FDA) has approved the use of one dose of 2009 H1N1 flu vaccine for persons 10 years of age and older.

In addition to protecting pregnant women from infection, infants less than 6 months old will not be able to be vaccinated so it is recommended that everyone who lives with or provides care for infants less than 6 months of age receive both the seasonal influenza vaccine and 2009 H1N1 influenza monovalent vaccine to provide protection for the infant. One recent study conducted in Bangladesh,

assessed the effectiveness of influenza immunization for mothers and their young infants. Inactivated influenza vaccine reduced proven influenza illness by 63% in infants up to 6 months of age. This study confirmed that maternal influenza immunization is a strategy with substantial benefits for both mothers and infants.

Q: Should the 2009 H1N1 influenza vaccine be given to someone who has had an influenza- like illness since between April and now? Do I need a test to know if I need the vaccine or not?

A. There is no test that can show whether a person had 2009 H1N1 influenza in the past. Many different infections, including influenza, can cause influenza-like symptoms such as cough, sore throat and fever. In addition, infection with one strain of influenza virus will not provide protection against other strains. People for whom influenza vaccine is recommended should receive the 2009 H1N1 vaccine, even if they had an influenza-like illness previously. It is not necessary to test a person who previously had an influenza-like illness. People for whom the 2009 H1N1 influenza vaccine is recommended should receive it, even if they have had an influenza-like illness previously, unless they can be certain they had 2009 H1N1 influenza based on a laboratory test that can specifically detect 2009 H1N1 viruses. CDC recommends that persons who were tested for 2009 H1N1 influenza discuss this issue with a healthcare provider to see if the test they had was either an RT-PCR or a viral culture that showed 2009 H1N1 influenza. There is no harm in being vaccinated if you had 2009 H1N1 influenza in the past.

Q: What are the possible side effects of the 2009 H1N1 influenza vaccine?

A. The side effects from 2009 H1N1 influenza vaccine are expected to be similar to those from seasonal flu vaccines. The most common side effects following vaccination are expected to be mild, such as soreness, redness, tenderness or swelling where the shot was given. Some people might experience headache, muscle aches, fever, nausea and fainting. If these problems occur, they usually begin soon after the shot and may last as long as 1-2 days. Like any medicines, vaccines can cause serious problems like severe allergic reactions. However life-threatening allergic reactions to vaccines are very rare. In 1976, an earlier type of swine flu vaccine was associated with cases of a severe paralytic illness called Guillain-Barre Syndrome (GBS) at a rate of approximately 1 case of GBS per 100,000 persons vaccinated. Some studies done since 1976 have shown a small risk of GBS in persons who received the seasonal influenza vaccine. This risk is estimated to be no more than 1 case of GBS per 1 million persons vaccinated. Since then, flu vaccines have not been clearly linked to GBS. GBS has a number of different causes, and GBS can occur in a person who has never received an influenza vaccine. The potential benefits of influenza vaccination in preventing serious illness, hospitalization, and death substantially outweigh these estimates of risk for vaccine-associated GBS.

Anyone who has a severe (life-threatening) allergy to eggs or to any other substance in the vaccine should not get the vaccine. People should always inform their immunization provider if they have any severe allergies, if they've ever had a severe allergic reaction following flu vaccination, or if they have ever had GBS.

Q. Can the family members of a pregnant woman receive the nasal spray vaccine?

A. Pregnant women should not receive the live nasal spray influenza vaccine but family and household members and other close contacts of pregnant women (including healthcare personnel) who are 2 through 49 years old, healthy* and not pregnant may receive live nasal spray vaccine.

Q. Can a pregnant healthcare worker administer the live nasal influenza vaccine?

A. Yes. No special precautions are (such as gloves) are necessary. Hands should be washed or cleaned with waterless hand sanitizer before and after administering the vaccine or having any direct contact with patients in a health care setting.